



Experience-based and evidence-oriented
Corticobasal® Implantology

Vol.18 EECI N° 3

December 2024

English Edition



4TH IF® CONSENSUS DOCUMENT ON STANDARD ALGORITHMS FOR OSSEOINTEGRATING AND OSSEOFIXATED IMPLANTS

IF® RESEARCH & EVIDENCE DEPARTMENT

ISSN 1864-1199 / e-ISSN 1864-1237

Published by IF Publishing, Germany
continued in 2024 as:

- **CMF Implant Direction (CMF, ID1)**
- **Journal of unwanted Results (JUR, ID2)**
- **Experience-based and evidence-oriented Corticobasal® Implantology (EECI, ID3)**



The Foundation of Knowledge

Editorial Board

Editor-in-Chief

Prof. Dr. Stefan Ihde
prof@ihde.com

Coordinating editor

Prof. Dr. Fadia Awadalkreem, Dubai
editor@implantfoundation.org

Editorial board (in alphabetic order)

Dr. Ashish Chakranarayan, India
Dr. Anita Doshi, India
Prof. Dr. Antonina Ihde, Belarus
Prof. Dr. Vitomir S. Konstantinovic, Serbia
Prof. Dr. Aleksandar Lazarov, Bulgaria
Dr. Juri Mitrushchenkov, Russia
Dr. Mahendra Perumal, India
Prof. Dr. Olga Sipić, Serbia
Prof. Dr. Jan Vares, Ukraine

Aspirants to the Editorial Board (in alphabetic order)

Dr. Salem Barmawi, Libya
Dr. Pablo Diaz, Ecuador
Dr. Marcos Daniel Gonzales, Colombia
Dr. Georg Huber, Germany
Dr. Mehul D Jani, India
Dr. Valeri Lysenko, Ukraine
Dr. Kiran Patel, India
Dr. Faizur Rahman, India
Dr. Egammal Sethuraman, India
Dr. Nikolai Spiridonov, Russia

Evidence reports and Critical Appraisals

IF® Research & Evidence Dept.

Annual Subscription

Euro 2.800

Copyright

Copyright © 2006 - 2024 by
International Implant Foundation
DE- 80802 Munich / Germany
www.implantfoundation.org

Contact

publishing@implantfoundation.org

CMF.Impl.dir.

ISSN 1864-1199
e-ISSN 1864-1237

Disclaimer

Hazards

Great care has been taken to maintain the accuracy of the information contained in this publication. However, the publisher and/or the distributor and/or the editors and/or the authors cannot be held responsible for errors or any consequences arising from the use of the information contained in this publication. The statements or opinions contained in editorials and articles in this publication are solely those of the authors thereof and not of the publisher, and/or the distributor, and/or the IIF.

The products, procedures and therapies described in this work are hazardous and are therefore only to be applied by certified and trained medical professionals in environment specially designed for such procedures. No suggested test or procedure should be carried out unless, in the user's professional judgment, its risk is justified. Whoever applies products, procedures and therapies shown or described in this publication will do this at their own risk. Because of rapid advances in the medical science, IF recommends that independent verification of diagnosis, therapies, drugs, dosages and operation methods should be made before any action is taken.

Although all advertising material which may be inserted into the work is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement by the publisher regarding quality or value of such product or of the claims made of it by its manufacturer.

Legal restrictions

This work was produced by IF Publishing, Munich, Germany. All rights reserved by IF Publishing. This publication including all parts thereof, is legally protected by copyright. Any use, exploitation or commercialization outside the narrow limits set forth by copyright legislation and the restrictions on use laid out below, without the publisher's consent, is illegal and liable to prosecution. This applies in particular to photostat reproduction, copying, scanning or duplication of any kind, translation, preparation of microfilms, electronic data processing, and storage such as making this publication available on Intranet or Internet.

Some of the products, names, instruments, treatments, logos, designs, etc. referred to in this publication are also protected by patents and trademarks or by other intellectual property protection laws (eg. «IF», «IIF» and the IF-Logo) are registered trademarks even though specific reference to this fact is not always made in the text.

Therefore, the appearance of a name, instrument, etc. without designation as proprietary is not to be construed as a representation by publisher that it is in the public domain.

Institutions' subscriptions allow to reproduce tables of content or prepare lists of Articles including abstracts for internal circulation within the institutions concerned. Permission of the publisher is required for all other derivative works, including compilations and translations. Permission of the publisher is required to store or use electronically any material contained in this journal, including any article or part of an article. For inquiries contact the publisher at the address indicated.



4TH IF[®] CONSENSUS DOCUMENT ON STANDARD ALGORITHMS FOR OSSEOINTEGRATING AND OSSEOFIXATED IMPLANTS

Authors

IF[®] Research & Evidence Department^{†*}

1. Evidence & Research Department, International Implant Foundation, Leopoldstr. 116, 80802 Munich, Germany

* Correspondence: prof@ihde.com

How to Cite This Article

IF[®] Research & Evidence Department
4th IF[®] Consensus Document on Standard Algorithms for Osseointegrating and Osseofixed Implants

Experience-based and evidence-oriented Corticobasal[®] Implantology (EECI), Vol. 18, No. 3, International Implant Foundation Publishing, 2024

1. Introduction

Osseointegration is defined as the direct contact between living bone and dental implant surface without interposed soft tissue at the light microscope level^[1, 2]. The nature of the bony contact zone depends on the type of bone that is in contact with the bone's surface. If osteonal bone is in direct contact with the implant surface, the outer layer of the osteons which show low / no mineralization are in contact with the bone. This was earlier misunderstood as „fibrointegration“.

Osseofixation is described as the anchorage of the implant by the surgeon on the cortical bone. This can be achieved through macro-mechanical anchorage on the first, second or third cortical, often followed by secondary osseointegration (or osseoadaptation) of those parts of the implant that are not in contact with a cortical or the bone at all^[3, 4]. This event depends on the functional stimulus on bone.

There are key differences between the algorithms for osseointegrating and osseofixed implants. These differences should be highlighted for educational purpose

to the treatment providers. Additionally, these differences should be considered during treatment planning, implant (anchoring) principles, surgical and prosthetic techniques, the overall probability of success, and follow-up measures. Note that the combination of implants for osseointegration and for osseofixation has never been scientifically proven.

The differences in algorithms include the following aspects:

1. The implantation principle
2. The patient selection
3. The surgical technique
4. The prosthetic approach
5. The follow-up and maintenance program
6. Implant failure modes

2. Differences Between the Algorithms for Osseointegrated and Osseofixed Implants

2.1 The Implantation Principle Osseointegrating Implants

- **Principle:** Osseointegrated implants rely on an ankylotic connection of the

endosseous implant surface with the cancellous bone. Osseointegration is the process by which the implant reaches direct contact with the bone, a contact that was not given right after placement of the implant. Osseointegration involves new bone formation and the growth of the bone towards the implant^[1, 2, 5, 6]. This process needs time, which is referred to as a „healing period“. However, this process was never scientifically proven in detail

- **Imaging:** Detailed 3D imaging (CBCT) is often used prior to implantation to accurately evaluate the quality and quantity of the bone, to plan the correct position of the implants, and to avoid neighboring vital anatomical structures. This is of great significance, especially in elderly patients, because the implants are often too large for the jawbone area in which they have to be placed from a static point of view^[7, 8]

Osseofixated Implants

- **Principle:** These implants anchor primarily in the second and third cortical of the jawbones through macro-mechanical anchorage, which provides high

stability^[8-10]. The implants can also pass through soft tissue (mucosa or muscle attachments in the area of the second or third cortex) or in / through cavities (enucleated cysts, maxillary sinus, nasal cavity). Third cortical anchorage refers to the anchorage in the cortical of the pterygoid process of the sphenoid bone

- **Imaging:** Panoramic X-rays are primarily used; however, CT scans or cone beam CT may be used postoperatively to verify the good anchorage of the implant in the second or third cortical. In cases of severe atrophy, CT scans or cone-beam CT may be used pre- and post-operatively. The pre-operative use of cone beam CT is to verify the possibilities of reaching the second cortical with the load transmitting implant parts and the drills. The quality of the second cortical is of less significance to be investigated. Hence, the functional loading of the bone will enhance mineralization in the shortest time. Always remember, „There is no possibility to improve the quality of the bone, except, by using it more and more for load transmission“. For weak (cortical) bone situations, the surgeon's decision should be to place more implants in the

affected jawbone or jaw segment, and not to try bone augmentation

2.2 The Patient Selection

Osseointegrating Implants

Patient selection is done very strictly and with various criteria for general health, bone quality, and quantity. Patients with insufficient bone usually require bone reconstruction procedures, a procedure that is always associated with additional risks^[11-13]. The medical condition of the patient, the oral hygiene, the patient's smoking habits, and the surgical experience of the treatment provider may limit this treatment option in many patients^[15-18].

Moreover, due to strict criteria for the minimal state of patient's health, this leads to frequent rejection of elderly patients, the group of patients who require implants most^[18]. From this point of view, osseointegrating (ankylosing) implants are mainly used in the age group between 25 and 60 years.

Osseofixed Implants

Osseofixed implants utilize the basal

bone for anchorage; hence, almost all the patients have sufficient jawbone for this straightforward and modern method of implantology. Bone augmentation is never part of the treatment plan if this method is used. The complete avoidance of any kind of bone manipulation increases the number of potential patients compared to all other methods of implantology.

Moreover, a high survival rate was reported in cases with a history of periodontitis and smoking^[19]. Nevertheless, treatment with intravenous bisphosphonates represents an incalculable risk for any bone surgery and logically, this also poses an increased risk with these implants^[9, 10].

2.3 The Surgical Technique

Osseointegrating Implants

- **Implant placement:** Osseointegrating implants are inserted into the cancellous bone after an implant osteotomy. This implant osteotomy (implant drilling) results in damaged osteonal systems along the osteotomy. In order to achieve primary stability the implant osteotomy is held smaller than the actual circumference of the implant. By inserting the implant, it is

pressed into the prepared bone cavity. This creates a primary bone-to-implant contact. Conical implants tend to achieve higher primary stability compared to cylindrical implants. According to the theory of „osseointegration“ this contact zone undergoes some remodeling over time, resulting in new bone formation referred to as a secondary bone contact^[5]. Thus, the stability of the implant relies mainly on the osseointegration process that takes place over time. Therefore, a healing period is necessary before the implant can be loaded. This dynamic process of bone remodeling directly impacts the primary stability of the implant. Thus, the stability of the implant may decrease during the first three to four weeks of „healing“^[6]. Any disturbance in this healing process can result in early implant failure. Literature related the cause of early implant failure (i.e.: before prosthesis insertion) to fibrous tissue formation between the implant and the surrounding bone in the early healing period^[20]. Despite the lack of a definitive definition for the primary predisposing factor leading to early implant failure, several factors may contribute, such as bone quality and quantity, the patient's medical

condition, smoking habit, the implant site and technique, the inserted implant size (diameter and length), the inserted torque, the surgical technique and skill, and their combination with the grafting procedure^[20–25]. Bone augmentation procedures are commonly used to add bone-like tissues to the available amount of (atrophied) natural bone. By applying such an „augmentation“, the skeleton of the patient is surgically modified to fit the osseointegrating implant^[24, 25]. Bone augmentations add costs and medical risks to the procedure. Additionally, they prolong the treatment due to the additionally required healing time for the „graft“^[26]. Many investigations^[27–29] have highlighted the association between early implant failure and bone grafting

- **Treatment steps:** While the placement of osseointegrating implants is a single-step procedure, additional surgical steps are necessary for these implants because their demand for bone is large. Only a few patients provide enough natural bone to host osseointegrating implants without bone augmentation

Osseofixated Implants

- **Implant placement:** Osseofixated implants may be placed into fresh extraction sockets or into healed bone sites long after the extraction. They are actively anchored by the surgeon directly in the second or third cortical. This creates immediate high stability^[9, 10, 30, 31]. While implants for osseointegration pass through the mucosa on the oral side of the alveolar crest, osseofixated implants penetrate also through the other side of the jaw bone (the second cortical) and often (depending on the anatomy of the site) also through or at least into the soft tissues on the other side of the second or third cortical. In some anatomic sites, the implants are reaching muscle attachment areas, which provides an exceptional strong protection against loss of mineralization due to the muscle forces^[31]
- **Treatment steps:** Immediate functional loading is the first method of choice. The prosthesis is connected rigidly connected to the implant within a period of 72 hours, i.e. before osteonal remodeling can set in. The prosthesis serves not only as masticatory device, but also as a (necessary) splint to stabilize the implants

2.4 The Prosthetic Treatment

Osseointegrating Implants

- **Loading protocol:** For the majority of the cases conventional delayed loading was emphasized; hence, the implant is only loaded after osseointegration is ensured, i.e., after several months. Immediate functional loading can be used in specific cases. The literature reported an increased incidence of implant failure with an immediate loading protocol in osseointegrating implants compared to delayed load^[32-35] and highlighted the role of smoking, implant length^[32], implant site^[33], and bone grafting
- **Design of the prosthesis:** It can be either fixed or removable; however, in the case of a denture-supported prosthesis, the final prosthesis is fitted after the healing phase

Osseofixated Implants

- **Loading protocol:** Due to stable anchorage in the second or third cortical bone layers, implants can be loaded immediately with high, predictable success and excellent biomechanical force distribution

- **Design of the prosthesis:** Fixed prostheses are used even in severely resorbed cases, with significant improvement in the patient's oral health and functions, as well as high reported patient satisfaction and improved quality of life^[19, 30, 36]
- **Dentures:** The dentures are attached within 72 hours after implant placement and they serve to stabilize the implants

Zirconium has been intensely used as a bridge material both on osseointegrating and osseofixed implants with high success^[37, 38].

2.5 The Follow-Up and Maintenance Program

Osseointegrating Implants

- **Follow-up:** Require regular check-ups to monitor the osseointegration and adjust the prostheses. The full healing process (i.e. the time period for adaptation and consolidation) takes up to two years. This means that after "osseointegration" is reached and the implant is loaded, more adaptation of the bone and changes in its morphology must be expected. Some of these changes are denomi-

nated as "periimplantitis". Although we know today that the onset of periimplantitis is a bone driven development and not mainly the result of an infection^[39, 40] regular oral hygiene is assumed necessary to avoid progression of periimplantitis. It is assumed that medical and local factors, smoking, and old age could be co-factors for the development of a periimplantitis^[14-18]

Osseofixed Implants

- **Follow-up:** Depending on the type of prosthetic construction and other circumstances of the treatment, the first control appointment will take place one to three months after the initial treatment, and then every nine to twelve months. After two to four years, most patients can be referred to longer check-up intervals. The aftercare protocol includes adjusting the chewing surfaces, increasing the height of the bite and the sagittal bite position, checking the implant stability, and restoring the free mobility of the bridge against the mucosa on the jawbone. Correcting early contacts and incorrect loading allows for the regression of any cortical overload osteolysis, provided

that this intervention occurs in a timely manner^[41]

The discussed differences in algorithm reflect the diverse principles of anchoring and treatment between osseointegrating and osseofixed implants and their understanding plays a significant for providing successful treatment results and assessing later the necessity of carrying out correct treatment planning, a successful surgical and prosthetic phase, the maintenance and the aftercare protocol in general (including corrective interventions).

2.6 Implant Failure Modes

Osseointegrating Implants

Literature has reported an association between the 2-stage rough implants and the incidence of periimplant mucositis or periimplantitis^[42, 43]. In a recent systematic review including 57 studies, Dreyer et al.^[42] reported an incidence range of 1.1% to 85.0% of periimplantitis at implant level, with a prevalence of 0.4% within three years to 43.9% within five years. Another study by Derks et al.^[43] showed that peri-

implantitis started early and that at years two and three, 52% and 66% of implants had bone loss of >0.5 mm, respectively. This complication could potentially impact the overall success of the implant and treatment, as it puts the entire implant at risk until it undergoes removal or exfoliation.

A consequence that significantly reduces the patient satisfaction and quality of life.

Furthermore, reports have documented a number of mechanical failures, including crown fractures, framework fractures, screw loosening, screw fractures, and fixture fractures^[44]. Hence, some implantologists may prefer the use of screw-type prostheses to ensure the possibility of unscrewing the prostheses to replace the broken part and / or failing to remove implants if mechanical complications occur.

Osseofixed Implants

Osseofixed implants never develop periimplantitis due to the smooth surface of the implant^[3, 9, 14], the only exception is if the treatment provider or the patient

roughens the polished implant surfaces.

On the other hand, osseofixed implants show the risk of mechanical overloading, especially during the first 24 months. Hence, to avoid this complication, regular clinical follow-up visits are mandatory to examine and adjust the patient's occlusion. The most important follow-up appointment is the three months' control^[3, 9, 41].

Moreover, these implants or the whole Bone-Implant-Prosthetic-System (BIPS) can fail if the stabilizing rigid splint of the BIPS has been completely or partially lost (prosthetic loosening of crowns; fractures of the bridge) or due to the use of temporary cement to fixate the prosthesis^[45].

3. Highlights on the Osseofixed Implants

A number of studies have shown that osseofixed implants provide a significant improvement in oral health compared to cases treated with the conventional osseointegrating implants. Many investigators documented a high survival and success rates, with a greater advantage of

lack of periimplantitis with osseofixed implants^[46–65].

Therefore, when planning to use conventional (osseointegrating) implants, it is crucial to ensure accurate patient information, particularly for patients with compromised ridge support that would require additional bone augmentation. For this reason, today updated requirements are valid. This includes information about the fact that in oral implantology bone augmentation is not necessary any more, and that the standard treatment protocol for osseofixed implants includes an immediate functional loading protocol.

The method of osseofixation also presents a breakthrough in the management of maxillofacial defects where these implants yield high long-term survival rates, significant improvement in aesthetics and function, the patient's self-esteem, patient satisfaction, and the quality of life^[60–65].

References

- 1** The Glossary of Implant Dentistry, The International Congress of Oral Implantologists ICOI, pp 92
- 2** Albeshri S, Greenstein G. Bone-to-Implant Contact: Its Relationship to Factors That Affect Primary and Biologic Implant Stability. Compendium of Continuing Education in Dentistry ; 2024; <https://cdeworld.com/courses/5416-bone-to-implant-contact-its-relationship-to-factors-that-affect-primary-and-biologic-implant-stability>
- 3** Ihde S. Indications and Treatment Modalities with Corticobasal Jaw Implants. Ann Maxillofac Surg. 2019 Jul-Dec;9(2):379-386. doi: 10.4103/ams.ams_142_19.
- 4** Ihde S, Ihde A, Lysenko V, Konstantinovic V, Palka L. New systematic terminology of cortical bone areas for osseo-fixated implants in strategic oral implantology. JJ Anatomy. 2016 Apr 11;1(2):7. <https://www.semanticscholar.org/paper/Systematic-Terminology-of-Cortical-Bone-Areas-for-Ihde-Ihde/18cb3994ed3eeda8999148d20772eedc5b46b196>
- 5** Cochran DL. The evidence for immediate loading of implants. J Evid Based Dent Pract. 2006 Jun;6(2):155-63. doi: 10.1016/j.jebdp.2006.04.018.
- 6** Barewal RM, Oates TW, Meredith N, Cochran DL. Resonance frequency measurement of implant stability in vivo on implants with a sand-blasted and acid-etched (SLA) surface. Int J Oral and Maxillofac Implants 2003;18(5):641-51.
- 7** Jacobs R, Salmon B, Codari M, Hassan B, Bornstein MM. Cone beam computed tomography in implant dentistry: recommendations for clinical use. BMC Oral Health. 2018 May 15;18(1):88. doi: 10.1186/s12903-018-0523-5.
- 8** Jacobs R, Quirynen M. Dental cone beam computed tomography: justification for use in planning oral implant placement. Periodontol 2000. 2014 Oct;66(1):203-13. doi: 10.1111/prd.12051. PMID: 25123769.
- 9** Ihde S. and Ihde A., Immediate Loading Guideline to Successful Implantology, 2010, International Implant Foundation, Munich, Germany.
- 10** Awadalkreem F, Khalifa N, Satti A, Suliman AM. Rehabilitation of Patients with Compromised Ridge Support Using Immediately Loaded Corticobasal Implant- supported Prostheses: A Prospective Observational Study. J Contemp Dent Pract. 2022 Oct 1;23(10):971-978. doi: 10.5005/jp-journals-10024-3416.
- 11** Pommer B, Hof M, Fädler A, Gahleitner A, Watzek G, Watzak G. Primary implant stability in the atrophic sinus floor of human cadaver maxillae: impact of residual ridge height, bone density, and implant diameter. Clin Oral Implants Res. 2014 Feb;25(2):e109-13. doi: 10.1111/clr.12071.
- 12** Spencer KR. Implant based rehabilitation options for the atrophic edentulous jaw. Australian dental journal. 2018 Mar;63:S100-7.

- 13** Flanagan D. Horizontal Alveolar Ridge Splitting and Expansion. *J Oral Implantol.* 2024 Jun 1;50(3):200-210. doi: 10.1563/aaid-joi-D-23-00186.
- 14** Carra MC, Blanc-Sylvestre N, Courtet A, Bouchard P. Primordial and primary prevention of peri-implant diseases: A systematic review and meta-analysis. *J Clin Periodontol.* 2023 Jun;50 Suppl 26:77-112. doi: 10.1111/jcpe.13790
- 15** Samara W, Moztarzadeh O, Hauer L, Babuska V. Dental Implant Placement in Medically Compromised Patients: A Literature Review. *Cureus.* 2024 Feb 14;16(2):e54199. doi: 10.7759/cureus.54199.
- 16** Chrcanovic BR, Albrektsson T, Wennerberg A. Periodontally compromised vs. periodontally healthy patients and dental implants: a systematic review and meta-analysis. *J Dent.* 2014 Dec;42(12):1509-27. doi: 10.1016/j.jdent.2014.09.013.
- 17** Ferreira SD, Martins CC, Amaral SA, Vieira TR, Albuquerque BN, Cota LOM, Esteves Lima RP, Costa FO. Periodontitis as a risk factor for peri-implantitis: Systematic review and meta-analysis of observational studies. *J Dent.* 2018 Dec;79:1-10. doi: 10.1016/j.jdent.2018.09.010.
- 18** Velasco-Ortega E, Carretero-Barrado L, Moreno-Muñoz J, Ortiz-García I, Núñez-Márquez E, Rondón-Romero JL, López-López J, Jiménez-Guerra Á, Monsalve-Guil L. Immediate Loading of Implants with Fixed Rehabilitations in Geriatric Edentulous Patients; Biological Complications. *J Clin Med.* 2023 Oct 16;12(20):6548. doi: 10.3390/jcm12206548.
- 19** Lazarov AB. The Impact of Diabetes, Smoking, and Periodontitis on Patients' Oral Health related Quality of Life after Treatment with Corticobasal Implants - An Evaluative Study. *Ann Maxillofac Surg.* 2021 Jul-Dec;11(2):253-260. doi: 10.4103/ams.ams_191_21.
- 20** Yari A, Fasih P, Alborzi S, Nikzad H, Romoozi E. Risk factors associated with early implant failure: A retrospective review. *Journal of Stomatology, Oral and Maxillofacial Surgery.* 2024 Sep 1;125(4):101749.
- 21** Krisam J, Ott L, Schmitz S, Klotz AL, Seyidaliyeva A, Rammelsberg P, Zenthöfer A. Factors affecting the early failure of implants placed in a dental practice with a specialization in implantology - a retrospective study. *BMC Oral Health.* 2019 Sep 5;19(1):208. doi: 10.1186/s12903-019-0900-8.
- 22** Palma-Carrió C, Maestre-Ferrín L, Peñarrocha-Oltra D, Peñarrocha-Diago MA, Peñarrocha-Diago M. Risk factors associated with early failure of dental implants. A literature review. *Med Oral Patol Oral Cir Bucal.* 2011 Jul 1;16(4):e514-7. doi: 10.4317/medoral.16.e514.
- 23** Sakka, S.; Baroudi, K.; Nassani, M.Z. Factors associated with early and late failure of dental implants. *J. Investig. Clin. Dent.* 2012, 3, 258–261. <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2041-1626.2012.00162.x>

- 24** Schwarz, F.; Ramanauskaite, A. It is all about peri-implant tissue health. *Periodontol 2000* 2022, 88, 9–12. <https://doi.org/10.1111/prd.12407>
- 25** Takamoli J, Pascual A, Martinez-Amargant J, Garcia-Mur B, Nart J, Valles C. Implant failure and associated risk indicators: A retrospective study. *Clin Oral Implants Res.* 2021 May;32(5):619-628. doi: 10.1111/clr.13732.
- 26** Sanz-Sánchez I, Sanz-Martín I, Ortiz-Vigón A, Molina A, Sanz M. Complications in bone-grafting procedures: Classification and management. *Periodontol 2000.* 2022 Feb;88(1):86-102. doi: 10.1111/prd.12413.
- 27** Munakata M, Kataoka Y, Yamaguchi K, Sanda M. Risk Factors for Early Implant Failure and Selection of Bone Grafting Materials for Various Bone Augmentation Procedures: A Narrative Review. *Bioengineering (Basel).* 2024 Feb 17;11(2):192. doi: 10.3390/bioengineering11020192.
- 28** Carr AB, Arwani N, Lohse CM, Gonzalez RLV, Muller OM, Salinas TJ. Early Implant Failure Associated With Patient Factors, Surgical Manipulations, and Systemic Conditions. *J Prosthodont.* 2019 Jul;28(6):623-633. doi: 10.1111/jopr.12978.
- 29** Antoun H, Karouni M, Abitbol J, Zouiten O, Jemt T. A retrospective study on 1592 consecutively performed operations in one private referral clinic. Part I: Early inflammation and early implant failures. *Clin Implant Dent Relat Res.* 2017 Jun;19(3):404-412. doi: 10.1111/cid.12477.
- 30** Awadalkreem F, Khalifa N, Satti A, Suliman AM. Rehabilitation of Patients with Compromised Ridge Support Using Immediately Loaded Corticobasal Implant- supported Prostheses: A Prospective Observational Study. *J Contemp Dent Pract.* 2022 Oct 1;23(10):971-978. doi: 10.5005/jp-journals-10024-3416.
- 31** Antonina I, Lazarov A, Gaur V, Lysenko V, Konstantinovic V, Grombkötö G, Palka L, Ihde S. Consensus Regarding 16 Recognized and Clinically Proven Methods and Sub- Methods for Placing Corticobasal® Oral Implants. *Ann Maxillofac Surg.* 2020 Jul- Dec;10(2):457-462. doi: 10.4103/ams.ams_62_20.
- 32** Prasant MC, Thukral R, Kumar S, Sadrani SM, Baxi H, Shah A. Assessment of Various Risk Factors for Success of Delayed and Immediate Loaded Dental Implants: A Retrospective Analysis. *J Contemp Dent Pract.* 2016 Oct 1;17(10):853-856. doi: 10.5005/jp-journals-10024-1943.
- 33** Chen J, Cai M, Yang J, Aldhohrah T, Wang Y. Immediate versus early or conventional loading dental implants with fixed prostheses: A systematic review and meta-analysis of randomized controlled clinical trials. *J Prosthet Dent.* 2019 Dec;122(6):516-536. doi: 10.1016/j.prosdent.2019.05.013.
- 34** Sanz-Sánchez I, Sanz-Martín I, Figuero E, Sanz M. Clinical efficacy of immediate implant loading protocols compared to conventional loading depending on the type of the restoration: a systematic review. *Clin Oral Implants Res.* 2015

Aug;26(8):964-982. doi: 10.1111/clr.12428.

35 Peñarrocha-Diago M, Demarchi CL, Maestre-Ferrín L, Carrillo C, Peñarrocha-Oltra D, Peñarrocha-Diago MA. A retrospective comparison of 1,022 implants: immediate versus nonimmediate. *Int J Oral Maxillofac Implants*. 2012 Mar-Apr;27(2):421-7. PMID: 22442783.

36 Awadalkreem F, Khalifa N, Satti A, Suleiman AM. The Influence of Immediately Loaded Basal Implant Treatment on Patient Satisfaction. *Int J Dent*. 2020 Apr 14;2020:6590202. doi: 10.1155/2020/6590202

37 Pozzi A, Arcuri L, Fabbri G, Singer G, Londono J. Long-term survival and success of zirconia screw-retained implant-supported prostheses for up to 12 years: A retrospective multicenter study. *J Prosthet Dent*. 2023 Jan;129(1):96-108. doi: 10.1016/j.prosdent.2021.04.026.

38 Matta RE, Eitner S, Stelzer SP, Reich S, Wichmann M, Berger L. Ten-year clinical performance of zirconia posterior fixed partial dentures. *J Oral Rehabil*. 2022 Jan;49(1):71-80. doi: 10.1111/joor.13276.

39 Ihde, S.; Ihde, A.; Sipic, O.; Patka, Ł. Peri-Implantitis: A New Definition Proposal Based on Unnatural Spatial Arrangement and Late Mechanical Coupling between Two Cortical Bone Layers during Osseointegration Phase. Part I. *Appl. Sci*. 2022,12,4317. <https://doi.org/10.3390/app12094317>

40 Ihde, S.; Ihde, A.; Sipic, O.; Patka, Ł. Peri-Implantitis: A New Definition Proposal Based on Unnatural Spatial Arrangement and Late Mechanical Coupling between Two Cortical Bone Layers during Osseointegration Phase Part II. *Appl. Sci*. 2022,12,5589. <https://doi.org/10.3390/app12115589>

41 Ihde S. and Ihde A. "Cookbook Mastication", 5th Edition, International Implant Foundation Publishing, Munich; 2021, ISBN 978-3-945889-29-9 xxxx.

42 Dreyer, H.; Grischke, J.; Tiede, C.; Eberhard, J.; Schweitzer, A.; Toikkanen, S.E.; Glöckner, S.; Krause, G.; Stiesch, M. Epidemiology and Risk Factors of Peri-Implantitis: A Systematic Review. *J. Periodontol Res*. 2018, 53, 657–681.

43 Derks, Schaller, et al Peri-implantitis - onset and pattern of progression. *J Clin Periodontol* 2016 Apr;43(4):383-8. doi: 10.1111/jcpe.12535. Epub 2016

44 Papaspyridakos P, Chen CJ, Chuang SK, Weber HP, Gallucci GO. A systematic review of biologic and technical complications with fixed implant rehabilitations for edentulous patients. *Int J Oral Maxillofac Implants*. 2012 Jan-Feb;27(1):102-10. PMID: 22299086.

45 Consensus No 8. Indications, Procedure and Application of Methods for Carrying Out Corrective Interventions with Corticobasal® Implants

46 Madhavan S, Jesudas PC, Afradh KM. Evaluation of Immediate Functional Loading with Single

Piece Implants. International Journal of Research and Review 2022 Vol. 9; Issue: 4:19-27.

47 Gosai H, Anchlia S, Patel K, Bhatt U, Chaudhari P, Garg N. Versatility of basal cortical screw implants with immediate functional loading. Journal of Maxillofacial and Oral Surgery. 2022 Sep;21(3):824-32.

48 Hosny AM. Evaluation of basal dental implant placement in basal bone of atrophic alveolar ridge. Al-Azhar Journal of Dental Science. 2022 Jan 1;25(1):71-8.

49 Abdali Y, Kashmoola S, Al-Mash'hadani M, Albaaj FS. A One-Year Retrospective Radiographic Assessment of Marginal Bone Loss Around Basal Implants and Impact of Multiple Risk Factors using Multivariate Analysis. Journal of Oral Research. 2021;10(5):1-8.

50 Lazarov A. A prospective cohort study of maxillary sinus complications in relation to treatments with Strategic Implants® penetrating into the sinus. Annals of Maxillofacial Surgery. 2020 Jul;10(2):365.

51 Awadalkreem F, Ahmad AG, Ihde S, Osman M. Effects of Corticobasal® implant protrusion inside the nasal and maxillary sinus. Annals of Maxillofacial Surgery. 2020 Jan;10(1):114.

52 Ihde S, Sipic O. Functional and Esthetic Indication for Dental Implant Treatment and Immediate Loading (2) Case Report and Considerations: Typical Attitudes of Dentists (and their Unions

toward Tooth Extractions and the Prevention of Early, Effective, and Helpful Dental Implant Treatment in the European Union. Ann Maxillofac Surg. 2019 Jul-Dec;9(2):470-474. doi: 10.4103/ams.ams_152_19.

53 Ihde S, Sipic O, Pałka L. A Single-Step Implant Treatment of a Patient Presenting with a High Smile Line (Gummy Smile) - Surgical, Prosthetic, and Financial Considerations of Full-Mouth Rehabilitation with One-Piece Implants Anchored in Cortical Bone - A Case Report. Ann Maxillofac Surg. 2020 Jul-Dec;10(2):512-517. doi: 10.4103/ams.ams_43_19.

54 Lazarov A. Soft-Tissue Augmentation in Periodontally Compromised Patients during Immediate Placement and Immediate Loading Dental Implant Surgery - A Retrospective Study. Ann Maxillofac Surg. 2023 Jan-Jun;13(1):37-43. doi: 10.4103/ams.ams_207_22.

55 Sahoo SK, Mishra S, Chinnannavar SN, Bajoria AA, Das A, Singh DK. Assessment of Oral Health-Related Quality of Life in Patients Receiving Corticobasal Dental Implants. J Pharm Bioallied Sci. 2023 Jul;15(Suppl 2):S1036-S1039. doi: 10.4103/jpbs.jpbs_236_23

56 Pałka ŁR, Lazarov A. Immediately Loaded Bicortical Implants Inserted in Fresh Extraction and Healed Sites in Patients with and Without a History of Periodontal Disease. Ann Maxillofac Surg. 2019 Jul-Dec;9(2):371-378

- 57** Debnath SC, Memmi L, Vishnuraj R, Bhagawati A. Assessment of peri implant bone loss in basal implant: a prospective clinical study. *International Journal of Oral and Maxillofacial Surgery*. 2019 May 1;48:55.
- 58** Oleg D, Alexander L, Konstantinović VS, Sipić O, Damir S, Miličić B. Immediate- functional loading concept with one-piece implants (beces/ becesn/kos/BOI®) in the mandible and maxilla- a multi-center retrospective clinical study. *Journal of Evolution of Medical & Dental Sciences/ JEMDS*. 2019;8(5):306-15.
- 59** Lazarov A. Immediate functional loading: Results for the concept of the Strategic Implant®. *Annals of Maxillofacial Surgery*. 2019 Jan;9(1):78.
- 60** Vitomir KS, Filip I, Vojkan L, Igor Đ, Lukasz P. Survival rate of disk and screw-type implants used for the retention of extraoral prostheses. *J Prosthet Dent*. 2022 Mar;127(3):499-507.
- 61** Awadalkreem F, Khalifa N, Ahmad AG, Suliman AM, Osman M. Prosthetic rehabilitation of maxillary and mandibular gunshot defects with fixed basal implant- supported prostheses: A 5-year follow-up case report. *Int J Surg Case Rep*. 2020;68:27-31. doi: 10.1016/j.ijscr.2020.02.025.
- 62** Gaur V, Doshi AG, Palka LR. Mandibular reconstruction using single piece zygomatic implant in conjunction with a reinforcing Fibular Graft Union: a case report. *International Journal of Surgery Case Reports*. 2020 Jan 1;73:347-54.
- 63** Awadalkreem F, Khalifa N, Ahmad AG, Suliman AM, Osman M. Oral rehabilitation of maxillofacial trauma using fixed corticobasal implant-supported prostheses: A case series. *Int J Surg Case Rep*. 2022 Nov;100:107769. doi: 10.1016/j.ijscr.2022.107769.
- 64** Gaur V, Perumal SM, Rahmaan F, Pałka Ł. A practical approach to orofacial rehabilitation in a patient after inferior maxillectomy and rhinectomy with mono framework construction supported on a zygomatic implant placed in the glabella: a case report. *Maxillofac Plast Reconstr Surg*. 2021 Jul 13;43(1):25. doi: 10.1186/s40902-021-00312-8.
- 65** Osman M, Ahmad AG, Awadalkreem F. A Novel Approach for Rehabilitation of a Subtotal Maxillectomy Patient with Immediately Loaded Basal Implant-Supported Prosthesis: 4 Years Follow-Up. *Case Rep Dent*. 2020 Feb 6; 2020:9650164. doi: 10.1155/2020/96501